



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Rebuttal from L. F. Barros and B. Weber

Barros, L F ; Weber, B

Abstract: Comment on CrossTalk opposing view: lack of evidence supporting an astrocyte-to-neuron lactate shuttle coupling neuronal activity to glucose utilisation in the brain. [J Physiol. 2018]

DOI: <https://doi.org/10.1113/jp275506>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-169164>

Journal Article

Published Version

Originally published at:

Barros, L F; Weber, B (2018). Rebuttal from L. F. Barros and B. Weber. Journal of Physiology, 596(3):355-356.

DOI: <https://doi.org/10.1113/jp275506>

CROSSTALK

Rebuttal from L. F. Barros and B. Weber

L. F. Barros¹ and B. Weber²¹Centro de Estudios Científicos, Valdivia 5110466, Chile²Institute of Pharmacology and Toxicology, University of Zurich, and Neuroscience Center Zurich, Zurich CH-8057, Switzerland

Email: fbarros@cecs.cl

Edited by: Francisco Sepúlveda & Ian Forsythe

Bak and Walls (2018) chose two specific aspects of brain metabolism: the usage of glucose by neurons and the differential expression of monocarboxylate transporters (MCTs) and lactate dehydrogenase (LDH) isoforms. We do not have strong objections to their more general claims.

The astrocyte-to-neuron lactate shuttle (ANLS) – right from the get-go – has not disputed neuronal glucose metabolism. We concur with our colleagues in that neurons have glucose transporters and that they metabolise glucose, as shown in NMDA-stimulated neurons in culture (Bak *et al.* 2009), and as reported by many others in diverse preparations. It is also well established that neurons have a conspicuously active Ca^{2+} -dependent malate–aspartate shuttle (Llorente-Folch *et al.* 2013), and are therefore well equipped to metabolise glucose and lactate at the same time during activity. The critical question about neuronal glucose consumption is not qualitative but quantitative. What we do not agree with is that there is preferential consumption of glucose by neurons as concluded by Patel *et al.* (2014) and Lundgaard *et al.* (2015). In our opinion, these two studies should be interpreted with caution for technical reasons that were advanced in our CrossTalk article.

We also agree with Bak and Walls (2018) in that differential expressions of MCTs and LDH in neurons and astrocytes do not demonstrate vectorial flux because the direction of flux is not governed by kinetics but by gradients. Indeed, compelling support for ANLS comes from the finding of steep astrocyte-to-neuron gradients for lactate in the somatosensory cortex of anaesthetised mice (Machler *et al.* 2016), and for NADH/NAD⁺ in

mouse hippocampal slices (Mongeon *et al.* 2016), the latter likely to remain steep during neuronal activity (Diaz-Garcia *et al.* 2017). In addition to these gradients the astrocytic membrane potential acts as a driving force, which extrudes lactate through ion channels (Sotelo-Hitschfeld *et al.* 2015; Karagiannis *et al.* 2016). Thus, the consumption of astrocytic lactate by neurons is not only favoured kinetically but also thermodynamically.

Beyond doubt anaesthesia affects metabolism, a caveat that also applies to non-physiological conditions prevalent during cell culture and tissue culture work. For these reasons, no single piece of data may be taken as definitive proof of ANLS. And it is true that many experimental findings should be replicated in awake behaving animals. What presently makes ANLS a strong hypothesis is the convergence of experimental evidence acquired by many groups using various models under diverse experimental conditions.

Call for comments

Readers are invited to give their views on this and the accompanying CrossTalk articles in this issue by submitting a brief (250 word) comment. Comments may be submitted up to 6 weeks after publication of the article, at which point the discussion will close and the CrossTalk authors will be invited to submit a 'LastWord'. Please email your comment, including a title and a declaration of interest, to jphysiol@physoc.org. Comments will be moderated and accepted comments will be published online only as 'supporting information' to the original debate articles once discussion has closed.

References

- Bak LK & Walls AB (2018). CrossTalk opposing view: Lack of evidence supporting an astrocyte-to-neuron lactate shuttle coupling neuronal activity to glucose utilisation in the brain. *J Physiol* **596**, 351–353.
- Bak LK, Walls AB, Schousboe A, Ring A, Sonnewald U & Waagepetersen HS (2009). Neuronal glucose but not lactate utilization is positively correlated with NMDA-induced neurotransmission and fluctuations in cytosolic Ca^{2+} levels. *J Neurochem* **109** (Suppl. 1), 87–93.
- Diaz-Garcia CM, Mongeon R, Lahmann C, Koveal D, Zucker H & Yellen G (2017). Neuronal stimulation triggers neuronal glycolysis and not lactate uptake. *Cell Metab* **26**, 361–374.

- Karagiannis A, Sylantyev S, Hadjihambi A, Hosford PS, Kasparov S & Gourine AV (2016). Hemichannel-mediated release of lactate. *J Cereb Blood Flow Metab* **36**, 1202–1211.
- Llorente-Folch I, Rueda CB, Amigo I, del Arco A, Saheki T, Pardo B & Satrustegui J (2013). Calcium-regulation of mitochondrial respiration maintains ATP homeostasis and requires ARALAR/AGC1-malate aspartate shuttle in intact cortical neurons. *J Neurosci* **33**, 13957–13971, 13971a.
- Lundgaard I, Li B, Xie L, Kang H, Sanggaard S, Haswell JD, Sun W, Goldman S, Blekrot S, Nielsen M, Takano T, Deane R & Nedergaard M (2015). Direct neuronal glucose uptake heralds activity-dependent increases in cerebral metabolism. *Nat Commun* **6**, 6807.
- Machler P, Wyss MT, Elsayed M, Stobart J, Gutierrez R, von Faber-Castell A, Kaelin V, Zuend M, San MA, Romero-Gomez I, Baeza-Lehnert F, Lengacher S, Schneider BL, Aebischer P, Magistretti PJ, Barros LF & Weber B (2016). In vivo evidence for a lactate gradient from astrocytes to neurons. *Cell Metab* **23**, 94–102.
- Mongeon R, Venkatachalam V & Yellen G (2016). Cytosolic NADH-NAD⁺ redox visualized in brain slices by two-photon fluorescence lifetime biosensor imaging. *Antioxid Redox Signal* **25**, 553–563.
- Patel AB, Lai JC, Chowdhury GM, Hyder F, Rothman DL, Shulman RG & Behar KL (2014). Direct evidence for activity-dependent glucose phosphorylation in neurons with implications for the astrocyte-to-neuron lactate shuttle. *Proc Natl Acad Sci USA* **111**, 5385–5390.
- Sotelo-Hitschfeld T, Niemeyer MI, Machler P, Ruminot I, Lerchundi R, Wyss MT, Stobart J, Fernandez-Moncada I, Valdebenito R, Garrido-Gerter P, Contreras-Baeza Y, Schneider BL, Aebischer P, Lengacher S, San Martin A, Le Douce J, Bonvento G, Magistretti PJ, Sepulveda FV, Weber B & Barros LF (2015). Channel-mediated lactate release by K^{+} -stimulated astrocytes. *J Neurosci* **35**, 4168–4178.

Additional information

Competing interests

None declared.

Author contributions

Both authors have contributed to the conception or design of the work, acquisition or analysis or interpretation of data for the work, and drafting the work or revising it critically for important intellectual content. Both authors have approved

the final version of the manuscript and agree to be accountable for all aspects of the work. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

Funding

L.F.B. is funded by Fondecyt Grant 1160317 and is supported by the Chilean Government

through the Centers of Excellence Basal Financing Program of CONICYT PB-01 granted to the Centro de Estudios Científicos (CECs). B.W. is supported by the University of Zurich, by the Swiss National Science Foundation, and the European Community Horizon 2020 GLINT project. He is a member of the Molecular Imaging Network Zurich.

Acknowledgements

We thank the members of the Barros and Weber labs for helpful discussions, and Karen Everett for critical reading of the manuscript.